



Reptiles



Reptile diversity information in J-BMP is based on species distribution data of all the reptiles natively found in Japan. We applied habitat suitability modeling on the distribution data of each species to predict their presences per each 1 km x 1 km grid cell across Japan. Once predicted their distributions, we integrated additional species-specific properties of endemism, extinction risk, and phylogenetic information to delineate geographic patterns of assorted aspects of biodiversity, such as taxonomic species richness, richness of endangered species and evolutionary distinctiveness of assemblages.



■ References ■

Isaac N.J.B., Turvey S.T., Collen B., Waterman C., Baillie J.E.M. (2007) Mammals on the edge: conservation priorities based on threat and phylogeny. PloS ONE DOI:10.1371/journal.pone.0000296. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0000296>

Databasing of occurrence observation

https://www.youtube.com/watch?time_continue=42&v=XaPFb_Gzvhs&feature=emb_logo

Guisan, A., Thuiller, W., and Zimmermann, N.E., (2017) Habitat suitability and distribution models with applications in R. Cambridge university press. DOI:10.1017/9781139028271 <https://www.cambridge.org/core/books/habitat-suitability-and-distribution-models/A17F74A3418DBF9ADA191A04C35187F9>

Lehtomäki J., Kusumoto B., Shiono T., Tanaka T., Kubota Y., Moilanen A. (2018) Spatial conservation prioritization for the East Asian islands: A balanced representation of multitaxon biogeography in a protected area network. Diversity and distributions, 2018: 25: 414–429 <https://onlinelibrary.wiley.com/doi/full/10.1111/ddi.12869>

